



PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Bc

Search 

for



Limits

Preview/Index

History

Clipboard

Details

About Entrez



Abstract

Show:

Sort



Text

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

LinkOut

Cubby

Related Resources

Order Documents

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

Privacy Policy

☐ 1: Genomics. 1997 Feb 15;40(1):55-62.

Related Articles, L

ELSEVIER SCIENCE
FULL-TEXT ARTICLE

Human neuroserpin (PI12): cDNA cloning and chromosomal localization to 3q26.

Schrimpf SP, Bleiker AJ, Brecevic L, Kozlov SV, Berger P, Osterwalder Krueger SR, Schinzel A, Sonderegger P.

Institute of Biochemistry, University of Zurich, Switzerland.

Neuroserpin is a novel serine protease inhibitor of the serpin family. It has been reported as a 55-kDa glycoprotein that is secreted from the axons of cultured central and peripheral nervous system neurons. In situ hybridization and Northern blot analyses at different developmental stages of the chicken revealed that neuroserpin is predominantly expressed in the nervous system and that most cells expressing neuroserpin can be qualified as bona fide neurons. We have isolated the full-length cDNA for human neuroserpin from a fetal retina cDNA library. The open reading frame of the cDNA of human neuroserpin, like that of its chicken counterpart, encodes a protein of 410 amino acids. The human and the chicken neuroserpin exhibit an amino acid sequence identity of 80%. Northern blot analysis of human organs demonstrated predominant expression of neuroserpin in the brain. By fluorescence in situ hybridization the human neuroserpin gene (HGMW-approved symbol PI12) was mapped to region q26 of chromosome 3.

PMID: 9070919 [PubMed - indexed for MEDLINE]



Abstract

Show:

Sort



Text

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)

Jul 17 2003 11:4

h

cb

hg

e

e

e

fcg

c

e

e

b

b

e



SCIENCE @ DIRECT

Register or Login: Password: Athens[Home](#)[Journals](#)[Abstract Databases](#)[Reference Works](#)[My Alerts](#)[My Profile](#)[Help](#)WELCOME GUEST USER [info](#)**Genomics**

Volume 40, Issue 1, 15 February 1997, Pages 55-62

doi:10.1006/geno.1996.4514 [?](#) Cite or link using doi
Copyright © 1997 Academic Press. All rights reserved.**This Document**► **Abstract**

- [Abstract + References](#)
- [PDF \(402 K\)](#)

Regular Article**Actions**

- [E-mail Article](#)

Human Neuroserpin (PI12): cDNA Cloning and Chromosomal Localization to 3q26^{*1}

Sabine P. Schrimpf, Andreas J. Bleiker, Lukrecija Brecevic, Serguei V. Kozlov, Philipp Berger, Thomas Osterwalder, Stefan R. Krueger, Albert Schinzel and Peter Sonderegger¹

Institute of Biochemistry, University of Zurich, CH-8057, Zurich, Switzerland

Institute of Medical Genetics, University of Zurich, CH-8001, Zurich, Switzerland

Received 15 July 1996; accepted 5 November 1996. Available online 18 April 2002.

Abstract

Neuroserpin is a novel serine protease inhibitor of the serpin family. It has been reported as a 55-kDa glycoprotein that is secreted from the axons of cultured central and peripheral nervous system neurons. *In situ* hybridization and Northern blot analyses at different developmental stages of the chicken revealed that neuroserpin is predominantly expressed in the nervous system and that most cells expressing neuroserpin can be qualified as *bona fide* neurons. We have isolated the full-length cDNA for human neuroserpin from a fetal retina cDNA library. The open reading frame of the cDNA of human neuroserpin, like that of its chicken counterpart, encodes a protein of 410 amino acids. The human and the chicken neuroserpin exhibit an amino acid sequence identity of 80%. Northern blot analysis of human organs demonstrated predominant expression of neuroserpin in the brain. By fluorescence *in situ* hybridization the human neuroserpin gene (HGMW-approved symbol PI12) was mapped to region q26 of chromosome 3.

^{*1} B. W. Festoff, Ed.

¹ To whom correspondence should be addressed at Institute of Biochemistry, University of Zurich, Winterthurerstrasse 190, CH-8057 Zurich, Switzerland. Telephone: +41-1-257 55 41. Fax: +41-1-363 79 47. E-mail: pson@bioc.unizh.ch.

Genomics

Volume 40, Issue 1 , 15 February 1997 , Pages 55-62

This Document

► **Abstract**

- [Abstract + References](#)
- [PDF \(402 K\)](#)

Actions

- [E-mail Article](#)

Home	Journals	Abstract Databases	Reference Works	My Alerts	My Profile	? Help
----------------------	--------------------------	------------------------------------	---------------------------------	---------------------------	----------------------------	------------------------

Send [feedback](#) to ScienceDirect

Software and compilation © 2003 ScienceDirect. All rights reserved.

ScienceDirect® is an Elsevier Science B.V. registered trademark.

Your use of this service is governed by [Terms and Conditions](#). Please review our [Privacy Policy](#) for details on how we protect information that you supply.